

- 4 (a) Let $U = \{p \in \mathcal{P}_4(\mathbf{R}) : p''(6) = 0\}$. Find a basis of U .
 (b) Extend the basis in (a) to a basis of $\mathcal{P}_4(\mathbf{R})$.
 (c) Find a subspace W of $\mathcal{P}_4(\mathbf{R})$ such that $\mathcal{P}_4(\mathbf{R}) = U \oplus W$.

$$\text{Let } U = \{p \in \mathcal{P}_4 \mid p''(6) = 0\}$$

Let $p \in \mathcal{P}_4$. Then,

$$p(x) = ax^4 + bx^3 + cx^2 + dx + e$$

$$p''(x) = 12ax^2 + 6bx + 2c$$

$$p''(6) = 12a(36) + 6b(6) + 2c = 0$$

$$0 = 432a + 36b + 2c$$

Assume that $a=1$ and $b=0$. Then

$$c = -216$$

$$\text{Let } p_1(x) = (x^4 - 216x^2)$$

Now assume that $a=0$ and $b=1$, then $c = -18$

$$\text{Let } p_2(x) = (x^3 - 18x^2)$$

Above rough work, we found two possible candidates for basis. Further, we have to two more. So, let

$$P_1(x) = 1 \Rightarrow P_1''(x) = 0 \Rightarrow P_1''(6) = 0 \text{ and}$$

$$P_2(x) = x \Rightarrow P_2''(x) = 0 \Rightarrow P_2''(6) = 0$$

All candidates are,

$$P_1(x) = 1, P_3(x) = (x^3 - 18x^2)$$

$$P_2(x) = x, P_4(x) = (x^4 - 216x^2)$$

Claim 1: P_1, P_2, P_3, P_4 are linearly independent.

Suppose that $\exists a_1, \dots, a_4 \in \mathbb{F}$ such that
$$a_1 + a_2x + a_3(x^3 - 18x^2) + a_4(x^4 - 216x^2) = 0$$

By comparing relevant terms

constant term $\rightarrow a_1 = 0$

x term $\rightarrow a_2 = 0$

x^3 term $\rightarrow a_3 = 0$

x^4 term $\rightarrow a_4 = 0$

Thus P_1, P_2, P_3, P_4 are linearly independent.

Claim 2 P_1, P_2, P_3, P_4 spans U .

Let $f(x) = ax^4 + bx^3 + cx^2 + dx + e \in U$.

$$= (ax^4 - 216ax^2) + (216ax^2 +$$

$$+ bx^3 - 18bx^2) + 18bx^2 + cx^2 + dx + e$$

$$= a(x^4 - 216x^2)$$

$$+ b(x^3 - 18x^2)$$

$$+ (216a + 18b)x^2 + cx^2$$

$$+ dx + e$$

$$= aP_4(x) + bP_3(x) + cP_2(x) + dP_1(x) + eP_0(x)$$

Therefore $P_1, \dots, P_4$ is a basis for U .

b)

Now let $P_1(x) = 1$

$$P_2(x) = x$$

$$P_3(x) = (x^3 - 18x^2)$$

$$P_4(x) = (x^4 - 216x^2)$$

$$P_5(x) = x^2$$

Claim 3: P_1, P_2, P_3, P_4, P_5 are linearly independent.

Suppose that $\exists a_1, \dots, a_5 \in \mathbb{F}$ such that

$$a_1 + a_2x + a_3(x^3 - 18x) + a_4(x^4 - 216x^2) + a_5x^5 = 0$$

By comparing relevant terms.

$$\text{constant term} \rightarrow a_1 = 0$$


$$x \text{ term} \rightarrow a_2 = 0$$

$$x^3 \text{ term} \rightarrow a_3 = 0$$

$$x^4 \text{ term} \rightarrow a_4 = 0$$

$$x^2 \text{ term} \rightarrow -18a_3 + 216a_4 + a_5 = 0$$
$$\Rightarrow a_5 = 0$$

Thus P_1, P_2, P_3, P_4, P_5 are linearly independent.



Claim 4: $P_1, \dots, P_5$ spans $\mathcal{P}_4$

$$\begin{aligned}\text{Let } g(x) &= k_4 x^4 + k_3 x^3 + k_2 x^2 + k_1 x + k_0 \\ &= k_4 x^4 - 216 k_1 x^2 + 216 k_1 x^2 \\ &\quad + k_3 x^3 - 18 k_3 x^2 + 18 k_3 x^2 \\ &\quad + k_2 x^2 + k_1 x + k_0 \\ &= k_4 (x^4 - 216 x^2) + k_3 (x^3 - 18 x^2) \\ &\quad + (216 k_1 + 18 k_3 + k_2) x^2 \\ &\quad + k_1 x + k_0\end{aligned}$$

Thus, $P_1, \dots, P_5$ span $\mathcal{P}_4$

Therefore, $P_1, \dots, P_5$ is a basis for $\mathcal{P}_4$

c)

$$\text{Let } W = \text{span}\{x^2\} = \{kx^2 \mid k \in \mathbb{R}\}$$

$$\text{Then } U + W = \mathcal{P}_4 \text{ (by (b))}$$

Further, It is easily see that $U \cap W = \{0\}$

$$\text{Therefore, } U \oplus W = \mathcal{P}_4$$